

RESEARCH ARTICLE SUMMARY

CANCER

The chromatin accessibility landscape of primary human cancers

M. Ryan Corces*, Jeffrey M. Granja*, Shadi Shams, Bryan H. Louie, Jose A. Seoane, Wanding Zhou, Tiago C. Silva, Clarice Groeneveld, Christopher K. Wong, Seung Woo Cho, Ansuman T. Satpathy, Maxwell R. Mumbach, Katherine A. Hoadley, A. Gordon Robertson, Nathan C. Sheffield, Ina Felau, Mauro A. A. Castro, Benjamin P. Berman, Louis M. Staudt, Jean C. Zenklusen, Peter W. Laird, Christina Curtis, The Cancer Genome Atlas Analysis Network, William J. Greenleaf†, Howard Y. Chang†

INTRODUCTION: Cancer is one of the leading causes of death worldwide. Although the 2% of the human genome that encodes proteins has been extensively studied, much remains to be learned about the noncoding genome and gene regulation in cancer. Genes are turned on and off in the proper cell types and cell states by transcription factor (TF) proteins acting on DNA regulatory elements that are scattered over the vast noncoding genome and exert long-range influences. The Cancer Genome Atlas (TCGA) is a global consortium that aims to accelerate the understanding of the molecular basis of cancer. TCGA has systematically collected DNA mutation, methyl-

ation, RNA expression, and other comprehensive datasets from primary human cancer tissue. TCGA has served as an invaluable resource for the identification of genomic aberrations, altered transcriptional networks, and cancer subtypes. Nonetheless, the gene regulatory landscapes of these tumors have largely been inferred through indirect means.

RATIONALE: A hallmark of active DNA regulatory elements is chromatin accessibility. Eukaryotic genomes are compacted in chromatin, a complex of DNA and proteins, and only the active regulatory elements are accessible by the cell's machinery such as TFs. The assay for

transposase-accessible chromatin using sequencing (ATAC-seq) quantifies DNA accessibility through the use of transposase enzymes that insert sequencing adapters at these accessible chromatin sites. ATAC-seq enables the genome-wide profiling of TF binding events that orchestrate gene expression programs and give a cell its identity.

RESULTS: We generated high-quality ATAC-seq data in 410 tumor samples from TCGA, identifying diverse regulatory landscapes across 23 cancer types. These chromatin accessibility profiles identify cancer- and tissue-specific DNA regulatory elements that enable classification of

ON OUR WEBSITE

Read the full article at <http://dx.doi.org/10.1126/science.aav1898>

tumor subtypes with newly recognized prognostic importance. We identify distinct TF activities in cancer based on differences in the inferred patterns of TF-DNA interaction and gene

expression. Genome-wide correlation of gene expression and chromatin accessibility predicts tens of thousands of putative interactions between distal regulatory elements and gene promoters, including key oncogenes and targets in cancer immunotherapy, such as *MYC*, *SRC*, *BCL2*, and *PDL1*. Moreover, these regulatory interactions inform known genetic risk loci linked to cancer predisposition, nominating biochemical mechanisms and target genes for many cancer-linked genetic variants. Lastly, integration with mutation profiling by whole-genome sequencing identifies cancer-relevant noncoding mutations that are associated with altered gene expression. A single-base mutation located 12 kilobases upstream of the *FGD4* gene, a regulator of the actin cytoskeleton, generates a putative de novo binding site for an NKX TF and is associated with an increase in chromatin accessibility and a concomitant increase in *FGD4* gene expression.


CONCLUSION: The accessible genome of primary human cancers provides a wealth of information on the susceptibility, mechanisms, prognosis, and potential therapeutic strategies of diverse cancer types. Prediction of interactions between DNA regulatory elements and gene promoters sets the stage for future integrative gene regulatory network analyses. The discovery of hundreds of noncoding somatic mutations that exhibit allele-specific regulatory effects suggests a pervasive mechanism for cancer cells to manipulate gene expression and increase cellular fitness. These data may serve as a foundational resource for the cancer research community. ■

The list of author affiliations is available in the full article online.

*These authors contributed equally to this work.

†Corresponding author. Email: howchang@stanford.edu (H.Y.C.); wjg@stanford.edu (W.J.G.)

Cite this article as M. R. Corces et al., *Science* 362, eaav1898 (2018). DOI: 10.1126/science.aav1898



Cancer gene regulatory landscape. Chromatin accessibility profiling of 23 human cancer types (left) in 410 tumor samples from TCGA revealed 562,709 DNA regulatory elements. The activity of these DNA elements organized cancer subtypes, identified TF proteins and regulatory elements controlling cancer gene expression, and suggested molecular mechanisms for cancer-associated inherited variants and somatic mutations in the noncoding genome. See main article for abbreviations of cancer types. Ref., reference; Var., variant.

RESEARCH ARTICLE

CANCER

The chromatin accessibility landscape of primary human cancers

M. Ryan Corces^{1*}, Jeffrey M. Granja^{1,2,3*}, Shadi Shams¹, Bryan H. Louie¹, Jose A. Seoane^{2,4,5}, Wanding Zhou⁶, Tiago C. Silva^{7,8}, Clarice Groeneveld⁹, Christopher K. Wong¹⁰, Seung Woo Cho¹, Ansuman T. Satpathy¹, Maxwell R. Mumbach^{1,2}, Katherine A. Hoadley¹¹, A. Gordon Robertson¹², Nathan C. Sheffield¹³, Ina Felau¹⁴, Mauro A. A. Castro⁹, Benjamin P. Berman⁷, Louis M. Staudt¹⁴, Jean C. Zenklusen¹⁴, Peter W. Laird⁶, Christina Curtis^{2,4,5}, The Cancer Genome Atlas Analysis Network[†], William J. Greenleaf^{1,2,3,15,16†}, Howard Y. Chang^{1,2,17,18†}

We present the genome-wide chromatin accessibility profiles of 410 tumor samples spanning 23 cancer types from The Cancer Genome Atlas (TCGA). We identify 562,709 transposase-accessible DNA elements that substantially extend the compendium of known cis-regulatory elements. Integration of ATAC-seq (the assay for transposase-accessible chromatin using sequencing) with TCGA multi-omic data identifies a large number of putative distal enhancers that distinguish molecular subtypes of cancers, uncovers specific driving transcription factors via protein-DNA footprints, and nominates long-range gene-regulatory interactions in cancer. These data reveal genetic risk loci of cancer predisposition as active DNA regulatory elements in cancer, identify gene-regulatory interactions underlying cancer immune evasion, and pinpoint noncoding mutations that drive enhancer activation and may affect patient survival. These results suggest a systematic approach to understanding the noncoding genome in cancer to advance diagnosis and therapy.

Cancer is a highly heterogeneous group of diseases, with each tumor type exhibiting distinct clinical features, patient outcomes, and therapeutic responses. The Cancer Genome Atlas (TCGA) was established to characterize this heterogeneity and understand the molecular underpinnings of cancer (1). Through large-scale genomic and molecular analyses, TCGA has revealed an exquisite diversity of genomic aberrations, altered transcriptional networks, and tumor subtypes that have engendered a more comprehensive understanding of disease etiologies and laid the foundations for new therapeutics and impactful clinical trials.

Work from TCGA and many others has demonstrated the importance of the epigenome to cancer initiation and progression (2). Profiling of cancer-specific coding mutations through whole-exome sequencing has identified prominent driver mutations in genes encoding chromatin remodel-

ing enzymes and modifiers of DNA methylation. These mutations drive alterations in the epigenome which, in turn, can establish the dysregulated cellular phenotypes that have become known as the hallmarks of cancer (3). Although many principles of chromatin regulation have been elucidated in cultured cancer cells, epigenomic studies of primary tumors are especially valuable, capturing the genuine ecosystem of heterotypic tumor and stromal cell interactions and the impacts of factors in the tumor microenvironment such as hypoxia, acidosis, and matrix stiffness (4). TCGA has carried out targeted DNA methylation profiling of more than 10,000 samples and, more recently, whole-genome bisulfite sequencing (WGBS) of 39 TCGA tumor samples (5). This data-rich resource has identified cancer-specific differentially methylated regions, providing an unprecedented view of epigenetic heterogeneity in cancer. In-

tegration of DNA methylation and additional TCGA data types has enabled the prediction of functional regulatory elements (6–8) and the identification of previously unknown cancer subtypes (9–13). Additional work has identified cancer-relevant variable enhancer loci by using histone modifications (14) and enhancer RNA sequencing (15). These studies represent, to date, the largest genome-wide epigenomic profiling efforts in primary human cancer samples.

Recently, the advent of the assay for transposase-accessible chromatin using sequencing (ATAC-seq) (16) has enabled the genome-wide profiling of chromatin accessibility in small quantities of frozen tissue (17). Because accessible chromatin is a hallmark of active DNA regulatory elements, ATAC-seq makes it possible to assess the gene regulatory landscape in primary human cancers. Combined with the richness of diverse, orthogonal data types in TCGA, the chromatin accessibility landscape in cancer provides a key link between inherited and somatic mutations, DNA methylation, long-range gene regulation, and, ultimately, gene expression changes that affect cancer prognosis and therapy.

Results

ATAC-seq in frozen human cancer samples is highly robust

We profiled the chromatin accessibility landscape for 23 types of primary human cancers, represented by 410 tumor samples derived from 404 donors from TCGA (protocol S1). These 23 cancer types are representative of the diversity of human cancers (Fig. 1A and data S1). From the 410 tumor samples, we generated technical replicates from 386 samples, yielding 796 genome-wide chromatin accessibility profiles (data S1). Given the size of this cohort, we first ensured that all generated ATAC-seq data could be uniquely mapped to the expected donor through comparison with single-nucleotide polymorphism (SNP) genotyping calls (fig. S1A). In all samples, the genotype from the ATAC-seq data generated in this study correlated most highly with previously published genotyping array data for the expected donor compared with that of all other 11,126 TCGA donors. All ATAC-seq data included in this study passed a minimum threshold of enrichment of signal over background (fig. S1, B to D, and data S1) with most samples showing a characteristic fragment size distribution with clear nucleosomal periodicity (fig. S1E). With this high-quality set of 410 tumor samples, we identified 562,709 reproducible (observed in more than one replicate) pan-cancer

¹Center for Personal Dynamic Regulomes, Stanford University, Stanford, CA 94305, USA. ²Department of Genetics, Stanford University School of Medicine, Stanford, CA 94305, USA. ³Program in Biophysics, Stanford University School of Medicine, Stanford, CA 94305, USA. ⁴Department of Medicine, Stanford University School of Medicine, Stanford, CA 94305, USA. ⁵Stanford Cancer Institute, Stanford University School of Medicine, Stanford, CA 94305, USA. ⁶Center for Epigenetics, Van Andel Research Institute, Grand Rapids, MI 49503, USA. ⁷Center for Bioinformatics and Functional Genomics, Cedars-Sinai Medical Center, Los Angeles, CA 90048, USA. ⁸Department of Genetics, Ribeirão Preto Medical School, University of São Paulo, Ribeirão Preto, SP CEP 14.040-905, Brazil. ⁹Bioinformatics and Systems Biology Laboratory, Polytechnic Center, Federal University of Paraná, Curitiba, PR CEP 80.060-000, Brazil. ¹⁰Department of Biomolecular Engineering, Center for Biomolecular Sciences and Engineering, University of California–Santa Cruz, Santa Cruz, CA 95064, USA. ¹¹Department of Genetics, Lineberger Comprehensive Cancer Center, The University of North Carolina at Chapel Hill, Chapel Hill, NC 27599, USA. ¹²Canada's Michael Smith Genome Sciences Center, BC Cancer Agency, Vancouver, BC V5Z 4S6, Canada. ¹³Center for Public Health Genomics, University of Virginia, Charlottesville, VA 22908, USA. ¹⁴National Cancer Institute, NIH, Bethesda, MD 20892, USA. ¹⁵Department of Applied Physics, Stanford University, Stanford, CA 94025, USA. ¹⁶Chan Zuckerberg Biohub, San Francisco, CA 94158, USA. ¹⁷Program in Epithelial Biology, Stanford University, Stanford, CA 94305, USA. ¹⁸Howard Hughes Medical Institute, Stanford University, Stanford, CA 94305, USA.

*The authors contributed equally to this work. †The Cancer Genome Atlas Analysis Network collaborators and affiliations are listed in the supplementary materials.

†Corresponding author. Email: howchang@stanford.edu (H.Y.C.); wjg@stanford.edu (W.J.G.)




Fig. 1. Pan-cancer ATAC-seq of TCGA samples identifies diverse regulatory landscapes. (A) Diagram of the 23 cancer types profiled in this study. Colors are kept consistent throughout figures. (B) Pan-cancer peak calls from ATAC-seq data. Peak calls from each cancer type are shown individually in addition to the 562,709 peaks that represent the pan-cancer merged peak set. Color indicates the type of genomic region overlapped by the peak. The numbers shown above each bar represent the number of samples profiled for each cancer type. UTR, untranslated region. (C) Overlap of cancer type-specific ATAC-seq peaks with Roadmap DNase-seq peaks from various tissues and cell types. Left: The percent of ATAC-seq peaks that are overlapped by one or more Roadmap peaks. Right: A heatmap of the percent overlap observed for each ATAC-seq peak set within the Roadmap DNase-seq peak set. Colors are scaled according to the minimum and maximum overlaps, which are indicated numerically to the right of the DNase-seq peak set names. The total number of ATAC-seq peaks (white to purple) or Roadmap DNase-seq regions (white to green) are shown colorimetrically. (D) Normalized ATAC-seq sequencing tracks of all 23 cancer types at the MYC locus. Each track represents the average accessibility per 100-bp bin across all

replicates. Known GWAS SNPs rs6983267 (COAD, PRAD) and rs35252396 (KIRC) are highlighted with light blue shading. Region shown represents chromosome 8 (chr8):126712193 to 128412193. (E) Normalized ATAC-seq sequencing tracks of five different COAD samples (top, orange) and KIRC samples (bottom, purple) shown across the same MYC locus as in Fig. 1D. Known GWAS SNPs rs6983267 (COAD, PRAD) and rs35252396 (KIRC) are highlighted with light blue shading. Region shown represents chr8:126712193 to 128412193. ACC, adrenocortical carcinoma; BLCA, bladder urothelial carcinoma; BRCA, breast invasive carcinoma; CESC, cervical squamous cell carcinoma; CHOL, cholangiocarcinoma; COAD, colon adenocarcinoma; ESCA, esophageal carcinoma; GBM, glioblastoma multiforme; HNSC, head and neck squamous cell carcinoma; KIRC, kidney renal clear cell carcinoma; KIRP, kidney renal papillary cell carcinoma; LGG, low grade glioma; LIHC, liver hepatocellular carcinoma; LUAD, lung adenocarcinoma; LUSC, lung squamous cell carcinoma; MESO, mesothelioma; PCPG, pheochromocytoma and paraganglioma; PRAD, prostate adenocarcinoma; SKCM, skin cutaneous melanoma; STAD, stomach adenocarcinoma; TGCT, testicular germ cell tumors; THCA, thyroid carcinoma; UCEC, uterine corpus endometrial carcinoma.

peaks of chromatin accessibility (Fig. 1B and data S2). These peaks were identified using a normalized peak score metric to enable direct comparison of peaks across samples of unequal sequencing depth, with each cancer type having an average of 105,585 peaks (range 56,125 to 215,978; Fig. 1B and fig. S1F; see methods). Reproducibility within the pan-cancer peak set was high for technical replicates (different nuclei from the same tumor sample; fig. S1, G and H), intratumor replicates (different samples from the same tumor; fig. S1I), and intertumor replicates (tumor samples from different donors; fig. S1, J and K).

Cancer chromatin accessibility extends the dictionary of DNA regulatory elements

The pan-cancer and cancer type-specific peak sets generated in this study enabled quantification of the number of DNA regulatory elements identified. To do this, we compared the regions defined by our pan-cancer and cancer type-specific peak sets to the regions defined by the Roadmap Epigenomics Project deoxyribonuclease I hypersensitive sites sequencing (DNase-seq) studies (18), finding a median of 34.4% overlap between the cancer type-specific peak sets and the various Roadmap tissue-type peak sets, with the strongest overlap occurring in the expected combinations (Fig. 1C and data S3). In total, about 65% of the pan-cancer peaks identified in this study had overlap with previously observed regulatory elements, highlighting both the consistency of our results with published datasets and the large number of additional putative regulatory elements observed in this study (Fig. 1C). Given the extensive coverage of Roadmap DNase-seq studies in healthy tissues, our results suggested that the disease context of cancer unveils the activity of additional DNA regulatory elements. Moreover, overlap of the ATAC-seq-defined DNA regulatory elements with chromatin immunoprecipitation sequencing (ChIP-seq)-defined ChromHMM regulatory states shows a strong enrichment of accessible chromatin sites in promoter and enhancer regions, as expected (fig. S1L). Although we profiled many samples in some cancer types [i.e., breast invasive carcinoma (BRCA), 75 tumor samples], we profiled fewer samples in multiple other cancer types (i.e., cervical squamous cell carcinoma, four tumor samples) (Fig. 1B). By estimating the number of unique peaks added with each additional sample, we found that cancer types have an estimated average of 169,822 total peaks (range 97,995 to 309,313) at saturation (fig. S1, M and N, and data S3), suggesting that profiling of additional samples of each cancer type would further expand the repertoire of regulatory elements.

Noncoding DNA elements reveal distinct cancer gene regulation and genetic risks

The *MYC* proto-oncogene locus provides a prime illustration of the diversity of the chromatin accessibility landscape across cancer types. *MYC* is embedded in a region with multiple DNA

regulatory elements and noncoding transcripts that regulate *MYC* in a tissue-specific fashion (19). We observed sufficient diversity in the chromatin accessibility landscape of the *MYC* locus to enable clustering of cancer types into two primary categories: (i) cancer types with extensive chromatin accessibility at 5' and 3' DNA elements, such as colon adenocarcinoma (COAD), and (ii) cancer types with chromatin accessibility primarily at 3' regulatory elements, such as kidney renal clear cell carcinoma (KIRC) (Fig. 1D). This trend is consistent across different samples of the same cancer type, as shown for COAD and KIRC (Fig. 1E) and is similar to the regulation observed in the *HOXD* locus (20).

Genome-wide association studies (GWAS) have identified numerous inherited risk loci for cancer susceptibility. However, many of these SNPs reside in the noncoding genome within known DNA regulatory elements. In the *MYC* locus, we identify known sites of chromatin accessibility, including peaks surrounding functionally validated GWAS cancer susceptibility SNPs (rs6983267 and rs35252396; Fig. 1, D and E). SNP rs6983267 is associated with increased susceptibility to colon adenocarcinoma and prostate adenocarcinoma (PRAD) (21–23), consistent with the presence of focal chromatin accessibility in these cancer types. However, SNP rs6983267 has not been previously associated with breast cancer or any squamous tumor types, which also have strong chromatin accessibility at this regulatory element in our ATAC-seq data (Fig. 1D). Similarly, SNP rs35252396 has been associated with KIRC and, in our data, shows strong accessibility in samples from kidney cancer types as well as breast and thyroid carcinoma, suggesting a potential role for these SNPs in previously unappreciated cancer contexts.

To visualize global patterns from our diverse ATAC-seq datasets, we performed Pearson correlation hierarchical clustering on distal and promoter elements (Fig. 2A). We found that distal elements exhibited a greater specificity and wider dynamic range of activity in association with cancer types, whereas promoter element accessibility was less cancer type-specific and showed similar patterns of correlation to global gene expression, as measured by RNA-seq (Fig. 2A). This functional specificity of distal regulatory elements was also previously observed in healthy tissues and in development (24, 25). Using *t*-distributed stochastic neighbor embedding (26) (t-SNE; Fig. 2B) and density clustering (27) (fig. S2A), we identified 18 distinct clusters, which we labeled based on the observed cancer-type enrichment (fig. S2B and data S3). We found strong concordance between this ATAC-seq-based clustering and the published multiomic iCluster scheme using TCGA mRNA-seq, microRNA (miRNA)-seq, DNA methylation, reverse-phase protein array (RPPA), and DNA copy number data (28) (Fig. 2, C and D). Comparing this clustering scheme to other TCGA-based clustering schemes, we observed the strongest concordance of our ATAC-seq clustering scheme with mRNA and cancer type (Fig. 2E).

This is consistent with the connection of chromatin accessibility to transcriptional output and the observation that ATAC-seq is strongly cell type-specific. Multiple observations can be made from these clusters: (i) Some cancer types split into two distinct clusters such as breast cancer (i.e., basal and nonbasal) and esophageal cancer (i.e., squamous and adenocarcinoma), (ii) cancer samples derived from the same tissue type often group together [i.e., kidney renal papillary cell carcinoma (KIRP) and KIRC], and (iii) some cancers group together across tissues as observed for squamous cell types (Fig. 3A and fig. S2B).

Cluster-specific regulatory landscapes identify patterns of transcription factor usage and DNA hypomethylation

Grouping of samples into defined clusters enables the determination of patterns in chromatin accessibility that are unique to each cluster. Using a framework that we term “distal binarization,” we identified the distal regulatory elements that are accessible only in a single cluster or small group of clusters (Fig. 3B, fig. S2C, and data S4). Of the 516,927 pan-cancer distal elements, 203,260 were found to be highly accessible in a single cluster or group of clusters (up to four clusters). These cluster-specific peak sets are enriched for motifs of transcription factors (TFs) with correlated gene expression that are known to be important for cancer and tissue identity (Fig. 3C, fig. S2D, and data S4). These include the androgen receptor (AR) in prostate cancer, forkhead box A1 (FOXA1) in nonbasal breast cancer, and melanogenesis-associated transcription factor (MITF) in melanoma. Moreover, these cluster-specific peak sets are enriched for known GWAS SNPs that are associated with cancers of the corresponding type (fig. S2E and data S5), highlighting that cancer-related GWAS SNPs tend to be located within or near cancer type-specific regulatory elements. The concordance of GWAS risk loci and cancer chromatin state has often been evaluated using cancer cell lines in the past, and our work provides a foundational map to evaluate noncoding GWAS SNPs in primary human cancers.

Consistent with published reports (12, 18, 29, 30), the degree of DNA methylation was anticorrelated with chromatin accessibility at regulatory elements, and regions lacking chromatin accessibility were more frequently methylated (fig. S2F). In particular, cluster-specific peak sets are hypomethylated in the relevant cancer types, though frequently methylated in other cancer types that lack accessibility in those peaks (fig. S2G). Consistent with these observations, which are based on DNA methylation array data, we see a strong depletion of DNA methylation at the center of both distal peaks and promoter peaks in a single patient profiled by WGBS (fig. S2H) (5). In our analysis of methylation levels within cluster-specific peak sets, we also identified a subgroup of brain cancers that exhibits DNA hypermethylation of peaks specific to nonbrain cancers (fig. S2G), likely caused by mutations in genes

that affect DNA methylation, such as isocitrate dehydrogenase 1 (*IDHI*) (fig. S3A). Similarly, we found that the subset of testicular germ cell tumors that are seminomas show a pattern of genome-wide DNA hypomethylation, consistent with a published report (31) (fig. S3B). Thus, a small number of TFs dominate the cis-regulatory landscape in each cancer type. These TFs are often the known key drivers of the respective

cancer or tissue type, and TF occupancy is associated with, and possibly causes, DNA hypomethylation of the corresponding DNA elements in cancer.


De novo identification of cancer subtypes from ATAC-seq data

Given the richness of the chromatin accessibility landscape, we explored the capacity of

ATAC-seq data to define molecular subtypes of cancer de novo. This analysis was limited to cancer types with sufficient available donors: BRCA (*N* = 74), PRAD (*N* = 26), and KIRP (*N* = 34). In KIRP, a gap statistic identified three distinct subgroups that are clearly separable by the first two principal components (Fig. 3D). The smallest of these subgroups contains four donors with very clear differences in ATAC-seq

Fig. 2. Chromatin accessibility profiles reveal distinct molecular subtypes of cancers.

(A) Pearson correlation heatmaps of ATAC-seq distal elements (left), ATAC-seq promoters (middle), and RNA-seq of all genes (right). Clustering orientation is dictated by the ATAC-seq distal element accessibility, and all other heatmaps use this same clustering orientation. Color scale values vary between heatmaps. Promoter peaks are defined as occurring between -1000 and +100 bp of a transcriptional start site. Distal peaks are all nonpromoter peaks. The total number of features (*N*) used for correlation is indicated above each Pearson correlation heatmap. (B) Unsupervised t-SNE on the top 50 principal components for the 250,000 most variable peaks across all cancer types. Each dot represents the merge of all technical replicates from a given sample. Color represents the cancer type shown above the plot. (C) Cluster residence heatmap showing the percent of each TCGA iCluster that overlaps with each ATAC-seq-based cluster. (D) ATAC-seq t-SNE clusters shown on the PanCanAtlas iCluster TumorMap. Each hexagon represents a cancer patient sample, and the positions of the hexagons are computed from the similarity of samples in the iCluster latent space. The color and larger size of the hexagon indicates the ATAC-seq cluster assignment. Samples that were not included in the ATAC-seq analysis are represented by smaller gray hexagons. The text labels indicate the cancer disease type. (E) Variation of information analysis of clustering schemes derived by using various data types from TCGA.



Downloaded from <http://science.sciencemag.org/> on October 27, 2018




Fig. 3. ATAC-seq clusters cancer samples to show cancer- and tissue-specific drivers. (A) Cluster residence heatmap showing the percent of samples from a given cancer type that reside within each of the 18 annotated ATAC-seq clusters. (B) Heatmap showing the ATAC-seq accessibility at distal elements ($N = 203,260$) identified to be cluster-specific by distal binarization. (C) Enrichment of TF motifs in peak sets identified in Fig. 3B. Enrichment is determined by a hypergeometric (HG) test $-\log_{10}(P \text{ value})$ of the motif's representation within the cluster-specific peaks compared to the pan-cancer peak set. Transcription factors shown represent a manually trimmed set of factors whose expression is highly correlated ($r > 0.4$) with the accessibility of the corresponding motif. Color represents the $-\log_{10}(P \text{ value})$ of the hypergeometric test. (D) Principal component analysis of the top 25,000 distal ATAC-seq peaks within the KIRP cohort ($N = 34$ samples). Each dot represents an individual sample. The color of the dots represents K -means clustering ($K = 3$ by gap statistic). (E) Distal binarization analysis based on the three K -means-defined groups identified and shown (by color) in

Fig. 3D. (F) Dot plot showing the number of nearby ATAC-seq peaks per gene from the group 1 distal binarization. Each dot represents a different gene. The MECOM gene (also called EVI1) is highlighted in red. (G) Normalized average sequencing tracks of K -means-defined groups 1, 2, and 3 at the MECOM locus. Peaks specific to group 1 are highlighted by light blue shading. (H) DNA copy number data at the MECOM locus in the three K -means-defined groups. Each dot represents an individual sample. CNV, copy number variation. (I) Average chromatin accessibility at peaks near the MECOM gene ($N = 42$ peaks) and RNA-seq gene expression of MECOM in KIRP samples ($N = 34$ samples). Each dot represents an individual donor. Dots are colored according to the clustering group colors shown in Fig. 3D. CPM, counts per million. (J) Kaplan-Meier analysis of overall survival of all KIRP donors in TCGA ($N = 287$) stratified by MECOM overexpressed ($N = 44$) and normal MECOM expression ($N = 243$). (K) Hazard plot of risk of dying from KIRP based on multiple covariates, including MECOM expression (hazard ratio = 5.2, 95% confidence interval = 2.4 to 11.0). Lines represent 95% confidence intervals.

accessibility identified by distal binarization (red coloring in Fig. 3E). Within the set of regulatory elements that are specific to this subgroup, we found 42 ATAC-seq peaks near the MDS1 and EVI1 complex locus (*MECOM*) gene (Fig. 3, F and G). Notably, the high chromatin accessibility of these *MECOM* peaks is not related to copy number amplification, as determined by DNA copy number array data (Fig. 3H). The expression of the *MECOM* gene is highly correlated with the mean ATAC-seq accessibility at these 42 ATAC-seq peaks [correlation coefficient (r) = 0.79, Fig. 3I]. Additionally, overexpression of *MECOM* is significantly associated with poorer overall survival across all available KIRP data from TCGA ($P = 2.2 \times 10^{-5}$, Cox proportional hazard test, Fig. 3J) with a hazard ratio of 5.2 (95% confidence interval = 2.4 to 11.0). This association is more substantial than lymph node status or patient age and is independent of cancer stage (Fig. 3K), indicating a potential prognostic role for these findings. Importantly, *MECOM* overexpression is not readily explained by any previously identified subgroups of KIRP, including subgroups with a CpG island methylator phenotype or mutations in the gene encoding fumarate hydratase, which have also been shown to confer poor overall survival (13). These results suggest that *MECOM* activation in KIRP identifies a previously unappreciated subgroup of patients with adverse outcomes, a finding that was uncovered by notable changes in the chromatin accessibility landscape of these samples.

Similarly, we found multiple distinct subgroups of PRAD and BRCA based on *K*-means clustering of the top 25,000 variable distal ATAC-seq peaks (fig. S3, C and D). In PRAD, these include subgroups driven by activity of AR, tumor protein P63 (*TP63*), and forkhead box-family TFs (fig. S3C). From an unsupervised analysis of breast cancer, we identified motifs of known TF drivers of luminal subtype identity, such as GATA binding protein 3 (*GATA3*) and FOXA1, as being enriched in the peak clusters specific to a subset of luminal samples (clusters 3 and 4, fig. S3D). We also identified a potential role for grainyhead-like (*GRHL*) TF motifs in basal breast cancer (32) (cluster 1, fig. S3D) and an overlapping role for nuclear factor I (*NFI*) in both basal and luminal A breast cancer (cluster 2, fig. S3D). Additionally, ATAC-seq data can be used to identify regions of copy number amplification de novo (33), enabling the classification of *HER2*-amplified cases of breast cancer (fig. S3, E to G).

Footprinting analysis defines TF activities in cancer

The high sequencing depth of the ATAC-seq data generated in this study (median of 56.7 million unique reads per technical replicate) enabled the profiling of TF occupancy at base-pair resolution through TF footprinting. TF binding to DNA protects the protein-DNA binding site from transposition while the displacement or depletion of one or more nucleosomes creates high DNA accessibility in the immediate flanking se-

quence. Collectively, these phenomena are referred to as the TF footprint. To characterize TF footprints, we adapted a recent approach (34) that quantifies the “flanking accessibility,” a measure of the accessibility of the DNA adjacent to a TF motif, and “footprint depth,” a measure of the relative protection of the motif site from transposition (Fig. 4A and data S6). To calculate these variables, we aggregated all insertions relative to the TF motif center, genome-wide (fig. S4A). To attempt to account for known Tn5 transposase insertion bias, we computed the hexamer frequency centered at Tn5 insertions and normalized for the expected bias at each position relative to the motif center (34) (see methods for potential limitations). Depending on the binding properties of a TF and its ability to affect local chromatin accessibility, changes in these properties would be detectable through this approach genome-wide (fig. S4, B and C). ChromVAR (35), a similar genome-wide approach which assesses the ability of a TF to affect flanking accessibility, identified a highly overlapping list of TFs (fig. S4D).

To uncover transcriptionally driven TF binding patterns, we correlated the RNA-seq gene expression of a given TF to its corresponding footprint depth and estimated flanking accessibility (data S6). A factor whose expression is sufficient to generate robust DNA binding would have a footprint depth and flanking accessibility that are significantly correlated to its gene expression [false discovery rate (FDR) < 0.1, purple dots in Fig. 4B], such as TP63 (Fig. 4, C and D) or NK2 homeobox 1 (*NKX2-1*) (Fig. 4, E and F). Increases in flanking accessibility and decreases in footprint depth are likewise accompanied by decreases in methylation (bottom of Fig. 4, D and F), consistent with the hypothesis that methylated DNA is less likely to be bound by TFs (36). Although footprint depth and flanking accessibility are often correlated, their divergence can suggest the modes of TF-DNA interaction. For example, factors whose expression is sufficient to cause opening of chromatin around the motif site but not to protect the motif site from transposition would be expected to only exhibit a significant correlation between gene expression and flanking accessibility (blue dots in Fig. 4B). This pattern of correlation could be caused by effects such as rapid TF off rates or low occupancy (fig. S4, E and F). Conversely, a small number of TFs have expression that is only significantly correlated with footprint depth (red dots in Fig. 4B). Though likewise rare, we also identified potential negative regulators whose expression is inversely correlated to gain of flanking accessibility and loss of footprint depth, such as the cut-like homeobox 1 (*CUX1*) TF (37) (Fig. 4B and fig. S4, G and H). This is the expected behavior of repressive TFs that bind DNA and lead to compaction of the neighboring sequence. These results predicted dozens of positive and negative regulators whose expression is strongly correlated with chromatin accessibility patterns near to their corresponding motif (fig. S4I and data S6). Overall, our

footprinting analysis identified putative TFs with activities correlated with gene expression.

Linking of DNA regulatory elements to genes predicts interactions relevant to cancer biology

The breadth and depth of this sequencing study enabled a robust association of ATAC-seq peaks with the genes that they are predicted to regulate. To do this, we implemented a strategy based on the correlation of ATAC-seq accessibility and gene expression across all samples (Fig. 5A, $N = 373$ with matched RNA-seq and ATAC-seq). Because promoter capture Hi-C data suggested that >75% of three-dimensional (3D) promoter-based interactions occur within a 500-kilobase pair (kbp) distance (38), we restricted the length scale of this analysis to 500 kbp to avoid spurious predictions. Using a conservative FDR cutoff of 0.01, we identified 81,323 unique links between distal ATAC-seq peaks and genes (Fig. 5B and data S7). Some of these links are driven by correlation across many cancer types (Fig. 5, C to E), whereas 70% are strongly driven by one cluster (Fig. 5F and data S7). To derive a final list of peak-to-gene links (Fig. 5B), putative links were filtered against (i) links whose correlation is strongly driven by DNA copy number amplification (“CNA”; fig. S5, A and B), (ii) regions with broad and high local correlation (“diffuse”; fig. S5, B and C), and (iii) links involving an ATAC-seq peak that overlaps the promoter of any gene (Fig. 5G). As expected, the histogram of distances between a peak and its target gene decays sharply with distance (39) (Fig. 5H). The expression of most genes is correlated with the activity of fewer than five different peaks (Fig. 5I), whereas most peaks are predicted to interact with a single gene (Fig. 5J). Additionally, this analysis found that only 24% of predicted links occur between an ATAC-seq peak and the nearest gene, indicating that the majority of predicted interactions skip over one or more genes and would not be possible to predict from primary sequence alone (Fig. 5K). In total, we predicted at least one peak-to-gene link for 8552 protein-coding genes, accounting for nearly half of all protein-coding genes in the human genome, including 48% of the curated Catalogue of Somatic Mutations in Cancer (COSMIC) cancer-relevant genes (data S7).

In addition to predicting peak-to-gene links across cancer types, we also predicted peak-to-gene links within breast cancer ($N = 74$ donors), identifying 9711 unique peak-to-gene links (fig. S5D and data S7). Of these links, 36% were also identified in our analysis of all cancer types (fig. S5E). Particularly important in these BRCA-specific links was the contribution of recurrent DNA CNA as a strong driver for spurious peak-to-gene correlation (Fig. 5G). These false-positive associations were removed through the use of published TCGA DNA copy number array data and a local correlation correction model, as mentioned above (see methods). The final predicted BRCA-specific links follow a similar distance

Fig. 4. Footprinting analysis identifies distinct TF activities in cancer. (A) Schematic illustrating the dynamics of TF binding (purple) and Tn5 insertion (green).


(B) Classification of TFs by the correlation of their RNA expression to the footprint depth and flanking accessibility of their motifs. Color represents whether the depth (red), flank (blue), or both (purple) are significantly correlated to TF expression below an FDR cutoff of 0.1. Each dot represents an individual deduplicated TF motif (see methods).

(C) TF footprinting of the TP63 motif (CIS-BP M2321_1.02) in lung cancer samples from the squamous (cluster 8) or adenocarcinoma (cluster 12) subtype. The Tn5 insertion bias track of TP63 motifs is shown below.

(D) Dot plots showing the footprint depth and flanking accessibility of TP63 motifs across all lung cancer samples studied. Each dot represents a unique sample. Color represents cancer type (top), RNA-seq gene expression (middle), or methylation beta value (bottom). Samples without matching RNA or methylation data are shown in gray.

(E) TF footprinting of the NKX2-1 motif (CIS-BP M6374_1.02) in lung cancer samples from the squamous (cluster 8) and adenocarcinoma (cluster 12) subtype. The Tn5 insertion bias track of NKX2-1 motifs is shown below.

(F) Dot plots showing the footprint depth and flanking accessibility of NKX2-1 motifs across all lung cancer samples studied. Each dot represents a unique sample. Color represents cancer type (top), RNA-seq gene expression (middle), or methylation beta value (bottom). Samples without matching RNA or methylation data are shown in gray.



distribution and peak-to-gene linking specificity as observed in the pan-cancer predicted links (fig. S5, F to I).

Many of these predicted peak-to-gene links occur in clusters where multiple nearby peaks are predicted to be linked to the same gene, indicating that these clusters of peak-to-gene links may function as part of a single regulatory unit or enhancer. Extending the width of the linked ATAC-seq peaks to 1500 bp allows for joining of these peaks into defined putative enhancer units (fig. S5J). This resulted in a total of 58,092 pan-cancer and 7622 BRCA-specific enhancer-to-gene links (data S7).

Validation and utility of predicted links between distal elements and genes

To verify a regulatory interaction for the predicted peak-to-gene links, we used a CRISPR interference (CRISPRi) (40) strategy using a catalytically dead Cas9 (dCas9) fused to a Kruppel-associated box (KRAB) domain, which mediates focal heterochromatin formation and functional silencing of noncoding DNA regulatory elements (Fig. 6A). In this way, targeting the distal peak region of a predicted peak-to-gene link would be expected to cause a decrease in the expression of the linked gene, located tens to hundreds of kilobases away. CRISPRi of a predicted distal regulatory element linked to *BCL2* (164 kbp, Fig. 5C) led to a significant reduction in *BCL2* gene expression in the luminal-like breast cancer MCF7 cell line but not in the basal-like MDA-MB-231 cell line (Fig. 6B), consistent with the role of *BCL2* as a luminal-specific survival factor (41). Similarly, CRISPRi of a distal regulatory element linked to the *SRC* oncogene (-49 kbp, Fig. 5D) led to a significant reduction in gene expression in both MCF7 cells and MDA-MB-231 cells (Fig. 6B). On a genome-wide scale, the predicted BRCA-specific peak-to-gene links show a strong enrichment in 3D chromosome conformation data from MDA-MB-231 cells (42), providing further support for our link prediction strategy (Fig. 6C). Moreover, we found that, of the peak-to-gene links predicted from BRCA ATAC-seq data that are also associated with a DNA methylation array CpG probe, 35% overlap with links predicted jointly from DNA methylation array and RNA-seq data in an ELMER analysis (8, 43) of the complete TCGA BRCA dataset ($N = 858$ tumors) ($P < 0.001$; Fig. 6D, fig. S6A, and data S8). These overlaps contain many luminal-specific and basal-specific links (fig. S6A), with a clear delineation between luminal (fig. S6B) and basal (fig. S6C) breast cancer samples. Integrating WGBS and ATAC-seq demonstrated the dynamics of methylation and chromatin accessibility and the overlap of predicted interactions at the non-basal *FOXA1* and basal forkhead box C1 (*FOXC1*) loci (fig. S6, D and E).

Similarly, previous work has leveraged TCGA RNA-seq data to infer transcriptional networks that consist of regulons, each of which is based on a TF regulator and its associated positive and negative target genes (fig. S7A) (44). For each

Downloaded from <http://science.sciencemag.org/> on October 27, 2018

regulon, every donor in the cohort can be assigned a positive, undefined, or negative regulon activity as measured by a differential enrichment score (dES) (45). Certain patterns of chromatin accessibility are expected on the basis of the target gene set and dES status of the donor (fig. S7B). For example, in donors with positive dES, chromatin at sites linked to positive target genes should be more accessible, whereas chromatin at sites linked to negative targets should be less accessible (fig. S7B). Examination of the estrogen receptor 1 (ESR1) regulon in the 74 BRCA donors profiled in this study identified 482 ATAC-seq distal peak-to-gene links corresponding to 124 ESR1 target genes (fig. S7C and data S8). Accessibility at these peaks is strongly concordant with expectations, further supporting the predicted links ($P < 1 \times 10^{-20}$, fig. S7D). Examination of this regulon across all TCGA

BRCA donors ($N = 1082$) showed a significant difference in overall survival between ESR1 dES-positive and -negative samples (fig. S7, E and F).

Together, pan-cancer and BRCA-specific peak-to-gene links further informed cancer-related GWAS polymorphisms, allowing the linkage of SNPs to putative gene targets with about 65% of all GWAS polymorphisms targeting a gene other than the closest gene on the linear genome (data S5). SNPs falling within peak-to-gene links were predicted to act on important cancer-related genes, including master regulators of cancer and tissue identity such as *NKX2-1* (fig. S7G) and *TP63* (fig. S7H). Focusing specifically on the BRCA peak-to-gene links for which published 3D chromosome conformation data are available, we found clear examples of GWAS SNPs interacting with distant, non-neighboring genes, such as *OSRI* (Fig. 6E and fig. S7I). More-

over, overlapping of the pan-cancer and breast cancer-specific peak-to-gene links with expression quantitative trait loci (eQTLs, where genetic variation at noncoding elements is associated with gene expression differences) from the Genotype-Tissue Expression (GTEx) project showed significant overlap in almost all comparisons ($N = 44$ of 48 comparisons) (fig. S7J and data S5). These results underscored our ability to use these predicted peak-to-gene links to generate key insights into published data and inform poorly understood aspects of cancer biology.

Identification of DNA regulatory elements related to immunological response to cancer

Of particular interest to current cancer therapy, immune infiltrates represent a substantial




Fig. 5. In silico linking of ATAC-seq peaks to genes. (A) Schematic of the in silico approach used to link ATAC-seq peaks in distal noncoding DNA elements to genes via correlation of chromatin accessibility and RNA expression. (B) Heatmap representation of the 81,323 pan-cancer peak-to-gene links predicted. Each row represents an individual link between one ATAC-seq peak and one gene. Color represents the relative ATAC-seq accessibility (left) or RNA-seq gene expression (right) for each link as a z-score. (C) Dot plot of the ATAC-seq accessibility and RNA-seq gene expression of a peak-to-gene link located 164 kbp away from the transcription start site of the *BCL2* gene (peak 498895) that is predicted to regulate its expression. Color represents the cancer type. Each dot represents an individual sample. (D) Same as in Fig. 5C but for a peak that is located 49 kbp away from the *SRC* gene (peak 525295). (E) Same

as in Fig. 5C but for a peak that is located 93 kbp away from the *PPARG* gene (peak 98874). (F) Same as in Fig. 5C but for a peak that is located 58 kbp away from the *ERBB3* gene (peak 381116). (G) Bar plot showing the number of predicted links that were filtered for various reasons. First, regions whose correlation is driven by DNA copy number amplification were excluded (“CNA”). Next, regions of high local correlation were filtered out (“Diffuse”). Lastly, peak-to-gene links where the peak overlapped a promoter region were excluded (“Promoter”). The remaining links (“Distal”) are used in downstream analyses. (H) Distribution of the distance of each peak to the transcription start site (TSS) of the linked gene. (I) Distribution of the number of peaks linked per gene. (J) Distribution of the number of genes linked per peak. (K) Distribution of the number of genes “skipped” by a peak to reach its predicted linked gene.

contribution to the overall tumor composition in solid tumors (46–48). We reasoned that infiltrating immune cells could contribute to our ATAC-seq data, both through actions on tumor cells and through increased chromatin accessibility at known immune-specific regulatory elements. Leveraging published ATAC-seq datasets from the human hematopoietic system (25) and data generated here from human dendritic cell subsets (Fig. 6F), we characterized each of our linked peaks by comparing its accessibility in immune cell types to its accessibility in bulk cancer samples (Fig. 6G). We reasoned that peaks that are more accessible in immune cells compared with our cancer cohort might be generated from immune cells associated with the tumor tissue (Fig. 6G). Additionally, we correlated each linked peak to the cytolytic activity score (49) of the tumor. The cytolytic activity score is based on the log-average gene expression of granzyme A and perforin 1, two CD8 T cell-specific markers. Linked peaks that exhibit high correlation to cytolytic activity might also be considered to be related to immune infiltration. Combining these two metrics, we identified peak-to-gene links expected to be highly relevant to immune infiltration, including links to genes relevant to antigen presentation and T cell response (Fig. 6H and data S9). The accessibility of these peak-to-gene links that were predicted to be immune-related is highly cor-

related with computationally predicted metrics of immune infiltration (46, 47) and inversely correlated with tumor purity (48) (Fig. 6I). One notable linked gene is programmed death ligand 1 (*PDL1*, also known as *CD274*), a key mediator of immune evasion by cancer and an important target for cancer immunotherapy. *PDL1* is linked to four putative distal regulatory elements that exhibit distinct chromatin accessibility across cancer types and are located as far as 43 kbp away from the *PDL1* transcription start site (Fig. 6, J and K). CRISPRi of each of these four putative *PDL1* regulatory elements significantly decreased, but did not abrogate, the expression of *PDL1* mRNA in at least one of the two breast cancer cell lines tested (MCF7 and MDA-MB-231 cells, Fig. 6L). These results support a model where the expression of *PDL1* is affected by the combined activity of multiple distal regulatory elements.

Identification of cancer-relevant noncoding mutations

In addition to identifying gene regulatory interactions in cancer, ATAC-seq combined with whole-genome sequencing (WGS) can be used to identify regulatory mutations driving cancer initiation and progression. For example, if a noncoding somatic mutation causes the generation of a TF binding site, this mutation could lead to an increase in chromatin accessibility


in cis and a concomitant increase in the observed frequency of the mutant allele in ATAC-seq as compared with that in WGS (Fig. 7A). Similarly, a mutation that inactivates a TF binding site can lead to a decrease in chromatin accessibility and a concomitant decrease in the observed frequency of the mutant allele. If such mutations in regulatory elements were to be functional in cancer, we might also expect that they increase or decrease chromatin accessibility beyond the expected distribution observed in nonmutated samples.

From the 404 donors profiled in this study, high-depth WGS data was available for 35 donors across 10 cancer types. These 35 donors had 374,705 called somatic mutations, with 32,696 falling within annotated ATAC-seq peaks and 2259 having at least 30 reads in both ATAC-seq and WGS data (data S10). Among these mutations were three separate occurrences of telomerase reverse transcriptase (*TERT*) gene promoter mutations (Fig. 7B), previously shown to generate de novo E26 transformation-specific (ETS) motif sites. ATAC-seq is especially well suited to identifying these *TERT* promoter mutations because the variant allele frequency is skewed owing to the increase in accessibility on the mutant allele (fig. S8A). Compared with the publicly available exome sequencing data from TCGA, where the *TERT* capture probes do not extend into the promoter region, ATAC-seq provided significantly

Fig. 6. Validation of long-range gene regulation of cancer in peak-to-gene links.

(A) Schematic of CRISPRi experiments performed. Each experiment uses three guide RNAs (gRNAs) to target an individual peak. The effect of this perturbation on the expression of the linked gene is determined with quantitative polymerase chain reaction (qPCR). (B) Gene expression changes by qPCR after CRISPRi of peaks predicted to be linked to the *BCL2* (peak 498895) and *SRC* (peak 525295) genes in MCF7 and MDA-MB-231 cells. Error bars represent the standard deviation of four technical replicates. *** $P < 0.001$ and NS is not significant by two-tailed Student's *t* test. (C) Meta-virtual circular chromosome conformation capture (4C) plot of predicted BRCA-specific peak-to-gene links with distances greater than 100 kbp. HiChIP interaction frequency is shown for the MDA-MB-231 basal breast cancer cell line as well as multiple populations of primary T cells. Th17, T helper 17 cell; Treg, regulatory T cell. (D) Bar plot showing the overlap of predicted ATAC-seq-based peak-to-gene links and DNA methylation-based ELMER predicted probe-to-gene links in BRCA, as a percentage of all ATAC-seq-based peak-to-gene links with a peak overlapping a methylation probe. The percentage of peak-to-gene links overlapping an ELMER probe-to-gene link (34.9%) is compared to the overlap with 1000 sets of randomized ELMER probe-to-gene links ($3.6 \pm 0.6\%$, $P \ll 0.001$). (E) Virtual 4C plot of the peak-to-gene link between rs4322801 and the *OSR1* gene. Normalized HiChIP interaction signal is shown for the MDA-MB-231 basal breast cancer cell line as well as multiple populations of primary T cells using the colors shown in Fig. 6C. ATAC-seq sequencing tracks are shown below for four BRCA samples and MDA-MB-231 cells with increasing levels of *OSR1* gene expression. The rs4322801 SNP (left) and *OSR1* gene (right) are highlighted by light blue shading. Region shown represents chr2:18999999 to 19425000. (F) Diagram of the hematopoietic differentiation hierarchy with differentiated cells colored as either B cells (green), T cell or natural killer (Tcell/NK) cells (blue), or myeloid cells (red). HSC, hematopoietic stem cell; LMPP, lymphoid-primed multipotent progenitor; CLP, common lymphoid progenitor; MPP, multipotent progenitor; CMP, common myeloid progenitor; GMP, granulocyte macrophage progenitor; HSPC, hematopoietic stem and progenitor cells; pDC, plasmacy-

toid dendritic cell; mDC, myeloid dendritic cell. (G) Schematic of the analysis shown in Fig. 6H. Peak-to-gene links are classified as related to immune infiltration if their accessibility is higher in immune cells than TCGA cancer samples and they are highly correlated to cytolytic activity. (H) Dot plot showing ATAC-seq peak-to-gene links with relevance to immune infiltration. Each dot represents an individual peak with a predicted gene link. Peaks that are related to immune cells have higher ATAC-seq accessibility in immune cell types compared to TCGA cancer samples. Peaks related to immune infiltration have a higher correlation to cytolytic activity. Color represents the cell type of the observation. The vertical dotted line represents the mean + 2.5 standard deviations above the mean for all ATAC-seq peak correlations to the cytolytic activity. The red shading indicates peak-to-gene links that are predicted to be related to immune infiltration. The blue shading indicates peak-to-gene links that are not predicted to be related to immune infiltration. NS, not significant. (I) Violin plots of the distribution of Spearman correlations across all peak-to-gene links predicted to be related to immune infiltration (red) or not (blue) with various metrics of tumor purity. (J) Normalized ATAC-seq sequencing tracks of the *PDL1* gene locus in six samples with variable levels of expression of the *PDL1* gene (right). Predicted links (red) are shown below for four peak-to-gene links (L1 to L4, peaks 293734, 293735, 293736, and 293740, respectively) to the promoter of *PDL1*. One of these peak-to-gene links (L2) overlaps an alternative start site for *PDL1* and was therefore labeled as a "promoter" peak during filtration. This peak-to-gene link was added to this analysis after manual observation. Region shown represents chr9:5400502 to 5500502. (K) Heatmap representation of the ATAC-seq chromatin accessibility of the 5000-bp region centered at each of the four peak-to-gene links shown in Fig. 6J. Each row represents a unique donor ($N = 373$) ranked by *PDL1* expression. The correlation of the chromatin accessibility of each peak with the expression of *PDL1* is shown below the plot. Color represents normalized accessibility. (L) Gene expression changes of the *PDL1* gene by qPCR after CRISPRi of peaks predicted to be linked to the *PDL1* gene in MCF7 and MDA-MB-231 cells. Error bars represent the standard deviation of four technical replicates. *** $P < 0.0001$ and ** $P < 0.05$ by two-tailed Student's *t* test.



higher sequencing coverage of the *TERT* promoter locus per read sequenced, enabling a more robust classification of *TERT* promoter mutations ($P < 1 \times 10^{-7}$, fig. S8B). Of the three *TERT* promoter mutations identified in the subset of donors with matched WGS, one mutation, in particular, leads to a significant increase in accessibility compared to the other nonmutated members of that cancer type (FDR < 0.0001, blue dot in Fig. 7, B and C). As expected, this increase in *TERT* promoter accessibility is associated with a concomitant increase in *TERT* gene expression (blue dot in Fig. 7C). *TERT* promoter mutations, however, are not the only way to increase *TERT* gene expression, because high *TERT* expression can also be observed in samples without identifiable *TERT* promoter mutations (Fig. 7C). Consistent with a previous report (50), differential motif analysis at the site of this *TERT* promoter mutation identified E74-like ETS tran-

scription factor 1 (ELF1) or ELF2 as the TF that likely binds to the de novo ETS motif (fig. S8C). In addition, we identified several mutations overlapping CCCTC-binding factor (CTCF) motif occurrences that are associated with decreased accessibility at that site (fig. S8, D and E). However, these mutations were relatively rare and often had only small effects on the accessibility of the CTCF motif site despite a known enrichment of somatic mutations in CTCF motif sites in cancer (51, 52). In addition to known *TERT* promoter mutations, integrative analysis of WGS and ATAC-seq data uncovered a mutation upstream of the FYVE RhoGEF and PH domain-containing 4 (*FGD4*) gene, a regulator of the actin cytoskeleton and cell shape. This mutation occurs in a bladder cancer sample where the variant allele frequency observed in ATAC-seq is markedly higher than the variant allele frequency observed in

WGS (Fig. 7B). This mutation is associated with a significant increase in accessibility compared to other bladder cancer samples in this cohort (Fig. 7, B and D) and is accompanied by a similar increase in *FGD4* mRNA (Fig. 7D). Moreover, this mutation upstream of the *FGD4* gene (referred to as eFGD4 for enhancer FGD4) leads to a level of accessibility that is higher than any of the other samples profiled by ATAC-seq in this study (fig. S8F) and a level of *FGD4* gene expression that is in the top 3% of all bladder cancer samples in TCGA (fig. S8G). As estimated by WGS data, this eFGD4 mutation is present in a subclone comprising about 13% of the tumor (Fig. 7E); however, the mutant allele is present in 96% of all ATAC-seq reads spanning this locus (Fig. 7E), demonstrating a strong preference for accessibility on the mutant allele. This eFGD4 mutation is analogous to, but potentially more potent than, the *TERT* promoter mutation




Fig. 7. Integration of WGS and ATAC-seq identifies cancer-relevant regulatory mutations.

(A) Schematic of how functional variants are identified in regulatory elements. The example shown depicts the *TERT* promoter. (B) Dot plot of the difference in variant allele frequency (VAF) of ATAC-seq and WGS and the changes in chromatin accessibility caused by the given variant with respect to other samples of the same cancer type. Variants with a higher variant allele frequency in ATAC-seq than WGS would be expected to cause an increase in accessibility. Each dot represents an individual somatic mutation. (C) Normalized ATAC-seq and RNA-seq of thyroid cancer donors profiled in this study. Each dot represents an individual donor. Blue dot represents the donor with a *TERT* promoter mutation shown in Fig. 7B. Other thyroid cancer donors known to harbor a *TERT* promoter mutation were excluded from this plot. The hinges of the box represent the 25th to 75th percentile. WT, wild type. (D) Normalized ATAC-seq and RNA-seq of bladder cancer donors profiled in this study. Each dot represents an individual donor. Purple dot represents the donor with a mutation upstream of the *FGD4* gene shown in Fig. 7B. The hinges of the box represent the 25th to 75th percentile. (E) Comparison of wild-type and mutant reads in WGS and ATAC-seq data at the *TERT* promoter and *FGD4* upstream region for the donors highlighted in (D) and (E). (F) Normalized ATAC-seq sequencing tracks of the *FGD4* locus in the 10 bladder cancer samples profiled in this study, including the one sample with a mutation predicted to generate a de novo NKX motif (TCGA-BL-A13J). Locus shown represents chr12:32335774 to 32435774. The mutation position is indicated by a black dotted line. The predicted enhancer region surrounding this mutation is highlighted by light blue shading. (G) Difference in motif score in the wild-type and mutant *FGD4* upstream region. Motif score represents the degree of similarity between the sequence of interest and the relevant motif. Each dot represents an individual motif. (H) Overlay of the NKX2-8 motif (CIS-BP M6377_1.02) and the wild-type and mutant sequences of the *FGD4* upstream region. (I) Kaplan-Meier survival analysis of TCGA bladder cancer patients with high (top 33%) and low (bottom 33%) expression for the *FGD4* gene.

described above (Fig. 7E). In the case of the eFGD4 mutation, this dramatic allele bias occurs because chromatin at this locus is not normally accessible in any of the bladder cancer samples profiled in this study (gray dots and tracks in Fig. 7, D and F) but becomes highly accessible in the context of the eFGD4 mutation (purple dot and track in Fig. 7, D and F). Differential motif analysis identified NKX factor motifs as the most strongly enriched in the sequence corresponding to the eFGD4 mutation (Fig. 7G), where a C-to-T transition at position two generated a perfect NKX2-8 motif de novo from a latent site (Fig. 7H). RNA-seq data from the mutated sample identified multiple expressed NKX TFs [transcripts per million (TPM) > 0.5], nominating NKX3-1, NKX2-3, and NKX2-5 as potential mediators of this DNA binding event (fig. S8H). From this, we hypothesized that the eFGD4 mutation creates a de novo binding site for an NKX TF which, upon binding to the DNA, leads to a broad increase in accessibility across the entire 12-kbp region upstream of the *FGD4* gene. This hypothesis was further supported by the observation that the ATAC-seq accessibility of the entire *FGD4* upstream locus occurs on a single phased allele (fig. S8I). Moreover, separation of subnucleosomal and nucleosome-spanning reads in the ATAC-seq data are consistent with protein binding at the site of the eFGD4 mutation (light blue shading in fig. S8I). Lastly, because higher *FGD4* expression is significantly associated with worse overall survival in bladder cancer (Fig. 7I and fig. S8J), this mutation could have functional consequence in this particular cancer. Whether the eFGD4 mutation or other enhancer mutations emerge as recurrent drivers of human cancer should be addressed in future studies. Our data identified multiple additional noncoding mutations associated with a concomitant gain of chromatin accessibility and increase in RNA expression (fig. S8, K to Q), and we anticipate that future work will uncover mechanisms underlying this type of regulatory mutation across all cancer types.

Discussion

Here we provide an initial characterization of the chromatin regulatory landscape in primary human cancers. This dataset identified hundreds of thousands of accessible DNA elements, expanding the dictionary of regulatory elements discovered through previous large-scale efforts such as The Roadmap Epigenomics Project. The identification of these additional elements was made possible through (i) our analysis of primary cancer specimens, (ii) greater saturation of some cancer and tissue types in our dataset, or (iii) potential differences between ATAC-seq and DNase-seq platforms. Nevertheless, the high overlap between the two datasets demonstrates the robustness of both platforms and the consistency of the observed results.

The exquisite cell type-specificity of distal regulatory elements from our ATAC-seq data enabled the classification of cancer types and the discovery of previously unappreciated cancer

subtypes. De novo clustering of TCGA samples based on chromatin accessibility strongly overlaps previous integrative clustering methods, identifying 18 distinct cancer clusters. Comparing this clustering scheme to other clustering schemes defined by cancer type, mRNA, miRNA, DNA methylation, RPPA, and DNA copy number alterations, we observed the strongest concordance of our clustering scheme with mRNA and cancer type, consistent with a close functional linkage between chromatin accessibility and transcriptional output. The strength of the observed associations is influenced by the features represented for each platform. For example, the DNA methylation clusters are based on cancer-specific promoter hypermethylation (28). Clustering based on DNA methylation at distal regulatory elements would likely show a stronger correlation with the ATAC-seq groupings, but distal regulatory element representation on the DNA methylation array used for these samples was too sparse to allow such an analysis. We also identified epigenetically distinct subtypes of kidney renal papillary cancer that have clear differences in overall survival. This cancer type-specific activity in DNA regulatory elements may arise via mutations within the regulatory element, pathologic transcription factor activity, or reflect the regulatory state of the tumor's cell of origin (e.g., stem cells). As the chromatin accessibility landscapes of additional primary cancer samples are profiled, we anticipate the identification of further epigenetic subdivisions with prognostic implications, potentially nominating avenues for therapeutic intervention.

The data generated in this study fully represents the cellular complexity of primary human tumors, comprising signals from tumor cells, infiltrating immune cells, stromal cells, and other normal cell types. In many ways, this complexity is advantageous because it allows complex systems-level analyses to be performed in the future, including cellular deconvolution approaches to understand the contributions of various cell types or cell states to the overall landscape of chromatin accessibility. However, the admixed nature of this signal also highlights the need for future work to profile the chromatin accessibility of matched healthy tissues to further refine the specific changes that drive cancer. Nevertheless, the chromatin accessibility profiles generated in this study represent the largest effort to date to characterize the regulatory landscape in primary human cancer cells.

Using this data-rich resource, we identified classes of TFs whose expression leads to different patterns in TF occupancy and motif protection. By integrating RNA-seq and ATAC-seq, we found factors whose expression is sufficient for both motif protection and nucleosome repositioning and demonstrated this binding to be inversely correlated with the level of DNA methylation at those binding sites. Despite this strong correlation, many sites of differential chromatin accessibility do not show differential methylation, demonstrating the complemen-

tarity of these two data types, perhaps owing to the presence of intermediate chromatin states such as poised promoters or enhancers (53, 54).

Moreover, integration of RNA-seq and ATAC-seq across the 373 donors with paired datasets enabled a quantitative model to link the accessibility of a regulatory element to the expression of predicted target genes. This workflow identified putative links for more than half of the protein-coding genes in the genome, informing the target genes of poorly understood GWAS SNPs and increasing our understanding of cancer gene regulatory networks. These predictions were further supported using 3D chromosome conformation data, and a subset were validated through CRISPRi experiments in breast cancer cell lines. However, profiling of chromosome conformation in primary cancer samples has not been performed on a large scale. Future work to produce maps of chromosome conformation in these or other primary cancer samples will improve our understanding of gene regulatory networks in cancer and further clarify the roles for certain GWAS-identified SNPs in cancer initiation and progression.

Lastly, through integration of WGS and ATAC-seq, we revealed a class of somatic mutations that occur in regulatory regions and lead to strong gains in chromatin accessibility. We demonstrated that these mutations likely lead to changes in nearby gene expression and affect genes whose expression is linked to poorer overall survival. Some of these mutations, such as those occurring in the *TERT* promoter, have been found to be recurrent whereas others, such as the mutation upstream of the *FGD4* gene, may be rare but functionally important. Because the enhancer functions are often distributed and latent enhancer sequences are pervasive in the genome, noncoding mutations in cancer may be especially challenging and require high-throughput functional assessment. Future larger-scale efforts to combine genome and epigenome sequencing will pave the way to tackling the noncoding genome in cancer.

Materials and methods summary

ATAC-seq data was generated from 410 tissue samples from the TCGA collection of primary human tumors. These samples spanned 23 different tumor types. These ATAC-seq data were used to cluster samples, identifying epigenetically defined patient subgroups. Moreover, TF regulators of cancer were defined, and footprinting of these regulators was correlated to gene expression to identify putative classes of TFs. A correlation-based model was developed to link ATAC-seq peaks to putative target genes. These putative links were validated using CRISPRi-based perturbation of the peak region followed by quantification of changes in gene expression. Publicly available HiChIP data and GTEx eQTL data were further used to support genome-wide peak-to-gene linkage predictions. Lastly, WGS and ATAC-seq were combined to identify noncoding mutations that affect chromatin accessibility in an allele-specific manner.

REFERENCES AND NOTES

- C. Hutter, J. C. Zenklusen, The Cancer Genome Atlas: Creating lasting value beyond its data. *Cell* **173**, 283–285 (2018). doi: [10.1016/j.cell.2018.03.042](https://doi.org/10.1016/j.cell.2018.03.042); pmid: 29625045
- W. A. Flavahan, E. Gaskell, B. E. Bernstein, Epigenetic plasticity and the hallmarks of cancer. *Science* **357**, eaal2380 (2017). doi: [10.1126/science.aal2380](https://doi.org/10.1126/science.aal2380); pmid: 28729483
- D. Hanahan, R. A. Weinberg, Hallmarks of cancer: The next generation. *Cell* **144**, 646–674 (2011). doi: [10.1016/j.cell.2011.02.013](https://doi.org/10.1016/j.cell.2011.02.013); pmid: 21376230
- M. Egeblad, E. S. Nakasone, Z. Werb, Tumors as organs: Complex tissues that interface with the entire organism. *Dev. Cell* **18**, 884–901 (2010). doi: [10.1016/j.devcel.2010.05.012](https://doi.org/10.1016/j.devcel.2010.05.012); pmid: 20627072
- W. Zhou et al., DNA methylation loss in late-replicating domains is linked to mitotic cell division. *Nat. Genet.* **50**, 591–602 (2018). doi: [10.1038/s41588-018-0073-4](https://doi.org/10.1038/s41588-018-0073-4); pmid: 29610480
- M. Almamun et al., Integrated methylome and transcriptome analysis reveals novel regulatory elements in pediatric acute lymphoblastic leukemia. *Epigenetics* **10**, 882–890 (2015). doi: [10.1080/15592294.2015.1078050](https://doi.org/10.1080/15592294.2015.1078050); pmid: 26308964
- Y. He et al., Improved regulatory element prediction based on tissue-specific local epigenomic signatures. *Proc. Natl. Acad. Sci. U.S.A.* **114**, E1633–E1640 (2017). doi: [10.1073/pnas.1618353114](https://doi.org/10.1073/pnas.1618353114); pmid: 28193886
- L. Yao, H. Shen, P. W. Laird, P. J. Farnham, B. P. Berman, Inferring regulatory element landscapes and transcription factor networks from cancer methylomes. *Genome Biol.* **16**, 105 (2015). doi: [10.1186/s13059-015-0668-3](https://doi.org/10.1186/s13059-015-0668-3); pmid: 25994056
- M. Ceccarelli et al., Molecular profiling reveals biologically discrete subsets and pathways of progression in diffuse glioma. *Cell* **164**, 550–563 (2016). doi: [10.1016/j.cell.2015.12.028](https://doi.org/10.1016/j.cell.2015.12.028); pmid: 26824661
- H. Noshmeh et al., Identification of a CpG island methylator phenotype that defines a distinct subgroup of glioma. *Cancer Cell* **17**, 510–522 (2010). doi: [10.1016/j.ccr.2010.03.017](https://doi.org/10.1016/j.ccr.2010.03.017); pmid: 20399149
- T. Hinoue et al., Genome-scale analysis of aberrant DNA methylation in colorectal cancer. *Genome Res.* **22**, 271–282 (2012). doi: [10.1101/gr.115723.110](https://doi.org/10.1101/gr.115723.110); pmid: 21659424
- P. A. Northcott et al., The whole-genome landscape of medulloblastoma subtypes. *Nature* **547**, 311–317 (2017). doi: [10.1038/nature22973](https://doi.org/10.1038/nature22973); pmid: 28726821
- The Cancer Genome Atlas Research Network, Comprehensive molecular characterization of papillary renal-cell carcinoma. *N. Engl. J. Med.* **374**, 135–145 (2016). doi: [10.1056/NEJMoa1505917](https://doi.org/10.1056/NEJMoa1505917); pmid: 26536169
- B. Akhtar-Zaidi et al., Epigenomic enhancer profiling defines a signature of colon cancer. *Science* **336**, 736–739 (2012). doi: [10.1126/science.1217277](https://doi.org/10.1126/science.1217277); pmid: 22499810
- H. Chen et al., A pan-cancer analysis of enhancer expression in nearly 9000 patient samples. *Cell* **173**, 386–399.e12 (2018). doi: [10.1016/j.cell.2018.03.027](https://doi.org/10.1016/j.cell.2018.03.027); pmid: 29625054
- J. D. Buenostro, P. G. Giresi, L. C. Zaba, H. Y. Chang, W. J. Greenleaf, Transposition of native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins and nucleosome position. *Nat. Methods* **10**, 1213–1218 (2013). doi: [10.1038/nmeth.2688](https://doi.org/10.1038/nmeth.2688); pmid: 24097267
- M. R. Corces et al., An improved ATAC-seq protocol reduces background and enables interrogation of frozen tissues. *Nat. Methods* **14**, 959–962 (2017). doi: [10.1038/nmeth.4396](https://doi.org/10.1038/nmeth.4396); pmid: 28846090
- A. Kundaje et al., Integrative analysis of 111 reference human epigenomes. *Nature* **518**, 317–330 (2015). doi: [10.1038/nature14248](https://doi.org/10.1038/nature14248); pmid: 25693563
- J. Schuijers et al., Transcriptional dysregulation of *MYC* reveals common enhancer-docking mechanism. *Cell Reports* **23**, 349–360 (2018). doi: [10.1016/j.celrep.2018.03.056](https://doi.org/10.1016/j.celrep.2018.03.056); pmid: 29641996
- G. Andrey et al., A switch between topological domains underlies *HoxD* genes collinearity in mouse limbs. *Science* **340**, 1234167 (2013). doi: [10.1126/science.1234167](https://doi.org/10.1126/science.1234167); pmid: 23744951
- M. Yeager et al., Genome-wide association study of prostate cancer identifies a second risk locus at 8q24. *Nat. Genet.* **39**, 645–649 (2007). doi: [10.1038/ng2022](https://doi.org/10.1038/ng2022); pmid: 17401363
- I. P. M. Tomlinson et al., A genome-wide association study identifies colorectal cancer susceptibility loci on chromosomes 10p14 and 8q23.3. *Nat. Genet.* **40**, 623–630 (2008). doi: [10.1038/ng.111](https://doi.org/10.1038/ng.111); pmid: 18372905

23. I. K. Sur *et al.*, Mice lacking a *Myc* enhancer that includes human SNP rs6983267 are resistant to intestinal tumors. *Science* **338**, 1360–1363 (2012). doi: [10.1126/science.1228606](https://doi.org/10.1126/science.1228606); pmid: [23118011](https://pubmed.ncbi.nlm.nih.gov/23118011/)
24. R. E. Thurman *et al.*, The accessible chromatin landscape of the human genome. *Nature* **449**, 75–82 (2012). doi: [10.1038/nature11232](https://doi.org/10.1038/nature11232); pmid: [22955617](https://pubmed.ncbi.nlm.nih.gov/22955617/)
25. M. R. Corces *et al.*, Lineage-specific and single-cell chromatin accessibility charts human hematopoiesis and leukemia evolution. *Nat. Genet.* **48**, 1193–1203 (2016). doi: [10.1038/ng.3646](https://doi.org/10.1038/ng.3646); pmid: [27526324](https://pubmed.ncbi.nlm.nih.gov/27526324/)
26. L. J. P. van der Maaten, G. E. Hinton, Visualizing data using t-SNE. *J. Mach. Learn. Res.* **9**, 2579–2605 (2008).
27. A. Rodriguez, A. Laio, Clustering by fast search and find of density peaks. *Science* **344**, 1492–1496 (2014). doi: [10.1126/science.1242072](https://doi.org/10.1126/science.1242072); pmid: [24970081](https://pubmed.ncbi.nlm.nih.gov/24970081/)
28. K. A. Hoadley *et al.*, Cell-of-origin patterns dominate the molecular classification of 10,000 tumors from 33 types of cancer. *Cell* **173**, 291–304.e6 (2018). doi: [10.1016/j.cell.2018.03.022](https://doi.org/10.1016/j.cell.2018.03.022); pmid: [29625048](https://pubmed.ncbi.nlm.nih.gov/29625048/)
29. M. Kulis *et al.*, Whole-genome fingerprint of the DNA methylome during human B cell differentiation. *Nat. Genet.* **47**, 746–756 (2015). doi: [10.1038/ng.3291](https://doi.org/10.1038/ng.3291); pmid: [26053498](https://pubmed.ncbi.nlm.nih.gov/26053498/)
30. H. S. Kim *et al.*, Pluripotency factors functionally premark cell-type-restricted enhancers in ES cells. *Nature* **556**, 510–514 (2018). doi: [10.1038/s41586-018-0048-8](https://doi.org/10.1038/s41586-018-0048-8); pmid: [29670286](https://pubmed.ncbi.nlm.nih.gov/29670286/)
31. H. Shen *et al.*, Integrated molecular characterization of testicular germ cell tumors. *Cell Reports* **23**, 3392–3406 (2018). doi: [10.1016/j.celrep.2018.05.039](https://doi.org/10.1016/j.celrep.2018.05.039); pmid: [29898407](https://pubmed.ncbi.nlm.nih.gov/29898407/)
32. S. Werner *et al.*, Dual roles of the transcription factor grainyhead-like 2 (GRHL2) in breast cancer. *J. Biol. Chem.* **288**, 22993–23008 (2013). doi: [10.1074/jbc.M113.456293](https://doi.org/10.1074/jbc.M113.456293); pmid: [23814079](https://pubmed.ncbi.nlm.nih.gov/23814079/)
33. S. K. Denny *et al.*, Nf1b promotes metastasis through a widespread increase in chromatin accessibility. *Cell* **166**, 328–342 (2016). doi: [10.1016/j.cell.2016.05.052](https://doi.org/10.1016/j.cell.2016.05.052); pmid: [27374332](https://pubmed.ncbi.nlm.nih.gov/27374332/)
34. S. Baek, I. Goldstein, G. L. Hager, Bivariate genomic fingerprinting detects changes in transcription factor activity. *Cell Reports* **19**, 1710–1722 (2017). doi: [10.1016/j.celrep.2017.05.003](https://doi.org/10.1016/j.celrep.2017.05.003); pmid: [28538187](https://pubmed.ncbi.nlm.nih.gov/28538187/)
35. A. N. Schep, B. Wu, J. D. Buenrostro, W. J. Greenleaf, ChromVAR: Inferring transcription-factor-associated accessibility from single-cell epigenomic data. *Nat. Methods* **14**, 975–978 (2017). doi: [10.1038/nmeth.4401](https://doi.org/10.1038/nmeth.4401); pmid: [28825706](https://pubmed.ncbi.nlm.nih.gov/28825706/)
36. Y. Yin *et al.*, Impact of cytosine methylation on DNA binding specificities of human transcription factors. *Science* **356**, eaaj2239 (2017). doi: [10.1126/science.aaj2239](https://doi.org/10.1126/science.aaj2239); pmid: [28473536](https://pubmed.ncbi.nlm.nih.gov/28473536/)
37. T. Ellis *et al.*, The transcriptional repressor CDP (Cut11) is essential for epithelial cell differentiation of the lung and the hair follicle. *Genes Dev.* **15**, 2307–2319 (2001). doi: [10.1101/gad.200101](https://doi.org/10.1101/gad.200101); pmid: [11544187](https://pubmed.ncbi.nlm.nih.gov/11544187/)
38. B. M. Javierre *et al.*, Lineage-specific genome architecture links enhancers and non-coding disease variants to target gene promoters. *Cell* **167**, 1369–1384.e19 (2016). doi: [10.1016/j.cell.2016.09.037](https://doi.org/10.1016/j.cell.2016.09.037); pmid: [27863249](https://pubmed.ncbi.nlm.nih.gov/27863249/)
39. C. P. Fulco *et al.*, Systematic mapping of functional enhancer-promoter connections with CRISPR interference. *Science* **354**, 769–773 (2016). doi: [10.1126/science.aag2445](https://doi.org/10.1126/science.aag2445); pmid: [27708057](https://pubmed.ncbi.nlm.nih.gov/27708057/)
40. L. S. Qi *et al.*, Repurposing CRISPR as an RNA-guided platform for sequence-specific control of gene expression. *Cell* **152**, 1173–1183 (2013). doi: [10.1016/j.cell.2013.02.022](https://doi.org/10.1016/j.cell.2013.02.022); pmid: [23452860](https://pubmed.ncbi.nlm.nih.gov/23452860/)
41. Y. H. Eom, H. S. Kim, A. Lee, B. J. Song, B. J. Chae, BCL2 as a subtype-specific prognostic marker for breast cancer. *J. Breast Cancer* **19**, 252–260 (2016). doi: [10.4048/jbc.2016.19.3.252](https://doi.org/10.4048/jbc.2016.19.3.252); pmid: [27721874](https://pubmed.ncbi.nlm.nih.gov/27721874/)
42. S. W. Cho *et al.*, Promoter of lncRNA gene *PVT1* is a tumor-suppressor DNA boundary element. *Cell* **173**, 1398–1412.e22 (2018). doi: [10.1016/j.cell.2018.03.068](https://doi.org/10.1016/j.cell.2018.03.068); pmid: [29731168](https://pubmed.ncbi.nlm.nih.gov/29731168/)
43. T. C. Silva, S. G. Coetzee, L. Yao, D. J. Hazelett, H. Noshmeh, B. P. Berman, Enhancer linking by methylation/expression relationships with the R package ELMER version 2. bioRxiv 148726 [Preprint], 11 June 2017. doi: [10.1101/148726](https://doi.org/10.1101/148726)
44. A. G. Robertson *et al.*, Comprehensive molecular characterization of muscle-invasive bladder cancer. *Cell* **171**, 540–556.e25 (2017). doi: [10.1016/j.cell.2017.09.007](https://doi.org/10.1016/j.cell.2017.09.007); pmid: [28988769](https://pubmed.ncbi.nlm.nih.gov/28988769/)
45. M. A. A. Castro *et al.*, Regulators of genetic risk of breast cancer identified by integrative network analysis. *Nat. Genet.* **48**, 12–21 (2016). doi: [10.1038/ng.3458](https://doi.org/10.1038/ng.3458); pmid: [26618344](https://pubmed.ncbi.nlm.nih.gov/26618344/)
46. V. Thorsson *et al.*, The immune landscape of cancer. *Immunity* **48**, 812–830.e14 (2018). doi: [10.1016/j.immuni.2018.03.023](https://doi.org/10.1016/j.immuni.2018.03.023); pmid: [29628290](https://pubmed.ncbi.nlm.nih.gov/29628290/)
47. K. Yoshihara *et al.*, Inferring tumour purity and stromal and immune cell admixture from expression data. *Nat. Commun.* **4**, 2612 (2013). doi: [10.1038/ncomms3612](https://doi.org/10.1038/ncomms3612); pmid: [24113773](https://pubmed.ncbi.nlm.nih.gov/24113773/)
48. S. L. Carter *et al.*, Absolute quantification of somatic DNA alterations in human cancer. *Nat. Biotechnol.* **30**, 413–421 (2012). doi: [10.1038/nbt.2203](https://doi.org/10.1038/nbt.2203); pmid: [22544022](https://pubmed.ncbi.nlm.nih.gov/22544022/)
49. M. S. Rooney, S. A. Shukla, C. J. Wu, G. Getz, N. Hacohen, Molecular and genetic properties of tumors associated with local immune cytolytic activity. *Cell* **160**, 48–61 (2015). doi: [10.1016/j.cell.2014.12.033](https://doi.org/10.1016/j.cell.2014.12.033); pmid: [25594174](https://pubmed.ncbi.nlm.nih.gov/25594174/)
50. M. M. Makowski *et al.*, An interaction proteomics survey of transcription factor binding at recurrent TERT promoter mutations. *Proteomics* **16**, 417–426 (2016). doi: [10.1002/pmic.201500327](https://doi.org/10.1002/pmic.201500327); pmid: [26553150](https://pubmed.ncbi.nlm.nih.gov/26553150/)
51. R. Katainen *et al.*, CTCF/cohesin-binding sites are frequently mutated in cancer. *Nat. Genet.* **47**, 818–821 (2015). doi: [10.1038/ng.3335](https://doi.org/10.1038/ng.3335); pmid: [26053496](https://pubmed.ncbi.nlm.nih.gov/26053496/)
52. D. Hnisz *et al.*, Activation of proto-oncogenes by disruption of chromosome neighborhoods. *Science* **351**, 1454–1458 (2016). doi: [10.1126/science.aad9024](https://doi.org/10.1126/science.aad9024); pmid: [26940867](https://pubmed.ncbi.nlm.nih.gov/26940867/)
53. R. D. Hawkins *et al.*, Distinct epigenomic landscapes of pluripotent and lineage-committed human cells. *Cell Stem Cell* **6**, 479–491 (2010). doi: [10.1016/j.stem.2010.03.018](https://doi.org/10.1016/j.stem.2010.03.018); pmid: [20452322](https://pubmed.ncbi.nlm.nih.gov/20452322/)
54. T. K. Kelly *et al.*, Genome-wide mapping of nucleosome positioning and DNA methylation within individual DNA molecules. *Genome Res.* **22**, 2497–2506 (2012). doi: [10.1101/gr.143008.112](https://doi.org/10.1101/gr.143008.112); pmid: [22960375](https://pubmed.ncbi.nlm.nih.gov/22960375/)

ACKNOWLEDGMENTS

We thank X. Ji and J. Collier for assistance in sequencing, P. Giresi and Epinomics for sharing advice and expertise related to ATAC-seq data analysis, and the members of the Greenleaf and Chang laboratories for thoughtful advice and critique.

Funding: Supported by the National Cancer Institute, NIH grants R35-CA209919 (to H.Y.C.), P50-HG007735 (to H.Y.C. and W.J.G.), and the Parker Institute for Cancer Immunotherapy (H.Y.C.). M.R.C. is supported by NIH K99-AG059918. Additional support through the NIH Genomic Data Analysis Networks 1U24CA210974-01 (J. Zhu), 1U24CA210949-01 (J. N. Weinstein), 1U24CA210978-01 (R. Beroukhim), 1U24CA210952-01 (S. J. Jones), 1U24CA210989-01 (O. Elemento), 1U24CA210990-01 (J. Stuart), 1U24CA210950-01 (R. Akbani), 1U24CA210969-01 (P.W.L.), and 1U24CA210988-01 (K.A.H.). W.J.G. is a Chan-Zuckerberg Biohub Investigator. H.Y.C. is an Investigator of the Howard Hughes Medical Institute. **Author contributions:** L.M.S., J.C.Z., W.J.G., and H.Y.C. conceived of and designed the project. M.R.C. and J.M.G. compiled figures and wrote the manuscript with the help of all authors. M.R.C. developed methodology for profiling frozen cancer tissues by ATAC-seq. S.S., B.H.L., and M.R.C. performed all tissue processing and ATAC-seq data generation. N.C.S. designed and wrote the ATAC-seq data processing pipeline with help from M.R.C. M.R.C. and J.M.G. processed all ATAC-seq data, and J.M.G.

performed all analyses and developed all analytical tools unless otherwise stated below. J.A.S. and C.G. performed survival analyses with supervision from C.C., A.G.R., and M.A.C. J.A.S. performed subtyping analysis for KIRP. W.Z. performed WGBS methylation analysis and variation of information clustering analysis with supervision from P.W.L. T.C.S. performed all ELMER analyses with supervision from B.P.B. C.G. performed all regulon analysis with supervision from A.G.R. and M.A.C. C.K.W. performed tumor map analysis. K.A.H. performed cluster coincidence analysis comparing ATAC-seq-derived clusters to TCGA iClusters. S.W.C. produced all Tn5 transposase used in this study and generated reagents and cell lines used in CRISPRi experiments. B.H.L., S.S., and M.R.C. performed CRISPRi experiments. A.T.S. generated human dendritic cell ATAC-seq data. J.M.G. and A.T.S. performed immune infiltration analysis. J.M.G. and M.R.M. performed HiChIP analysis. M.R.C. performed all analysis for identifying noncoding mutations from WGS and ATAC-seq data. I.F. coordinated all TCGA analysis working group efforts. J.C.Z. selected tumor samples to profile in this study. P.W.L. and W.J.G. co-chaired the TCGA analysis working group. C.C. provided expertise relevant to pan-cancer data analysis. H.Y.C. and W.J.G. supervised overall data generation and analysis. All authors listed under “The Cancer Genome Atlas Analysis Network” provided valuable input and expertise. **Competing interests:** H.Y.C. is a co-founder of Accent Therapeutics and Epinomics and is an adviser of 10X Genomics and Spring Discovery. W.J.G. is a co-founder of Epinomics and an adviser to 10X Genomics, Guardant Health, CentriLion, and NuGen. C.C. is an adviser to GRAIL. Stanford University holds a patent on ATAC-seq, on which H.Y.C. and W.J.G. are named as inventors. **Data and materials availability:** Processed data not provided in the supplementary data files are available through our TCGA Publication Page (<https://gdc.cancer.gov/about-data/publications/ATACseq-AWG>). This includes pan-cancer raw and normalized counts matrices, cancer type-specific peak calls, cancer type-specific raw and normalized count matrices, and bigWig track files for all technical replicates. Raw ATAC-seq data as fastq or aligned BAM files will be made available through the NIH Genomic Data Commons portal (<https://portal.gdc.cancer.gov/>). ATAC-seq data corresponding to human plasmacytoid dendritic cells and myeloid dendritic cells (the only non-TCGA data generated here) is available through SRA BioProject PRJNA491478. The ATAC-seq peak accessibility and computed peak-to-gene linkage predictions are publicly available for interactive visualization and exploration at the UCSC Xena Browser (<https://atascseq.xenahubs.net>). Sample-level ATAC-seq data across all 404 donors assayed can be visualized side-by-side with all other data from TCGA, including gene expression, DNA methylation from both Illumina 450K array and WGBS platforms, and ELMER enhancer analysis results, as well as the latest survival data and mutation calls from the Genomic Data Commons. ATAC-seq data can be queried by gene, genomic position, or individual peaks. The UCSC Xena Browser makes this rich resource available for interactive online analysis and visualization by the larger scientific community. Samples from the TCGA project can only be used for TCGA efforts owing to restrictions in the material transfer agreement used for acquisition. No external groups can access the tissue or analytes.

SUPPLEMENTARY MATERIALS

www.sciencemag.org/content/362/6413/eaav1898/suppl/DC1
TCGA Analysis Network Collaborators
Materials and Methods
Protocol S1
Figs. S1 to S8
References (55–78)
Data S1 to S10

22 August 2018; accepted 28 September 2018
10.1126/science.aav1898

The chromatin accessibility landscape of primary human cancers

M. Ryan Corces, Jeffrey M. Granja, Shadi Shams, Bryan H. Louie, Jose A. Seoane, Wanding Zhou, Tiago C. Silva, Clarice Groeneveld, Christopher K. Wong, Seung Woo Cho, Ansuman T. Satpathy, Maxwell R. Mumbach, Katherine A. Hoadley, A. Gordon Robertson, Nathan C. Sheffield, Ina Felau, Mauro A. A. Castro, Benjamin P. Berman, Louis M. Staudt, Jean C. Zenklusen, Peter W. Laird, Christina Curtis, The Cancer Genome Atlas Analysis Network, William J. Greenleaf and Howard Y. Chang

Science **362** (6413), eaav1898.
DOI: 10.1126/science.aav1898

Cancer chromatin accessibility landscape

The Cancer Genome Atlas (TCGA) provides a high-quality resource of molecular data on a large variety of human cancers. Corces *et al.* used a recently modified assay to profile chromatin accessibility to determine the accessible chromatin landscape in 410 TCGA samples from 23 cancer types (see the Perspective by Taipale). When the data were integrated with other omics data available for the same tumor samples, inherited risk loci for cancer predisposition were revealed, transcription factors and enhancers driving molecular subtypes of cancer with patient survival differences were identified, and noncoding mutations associated with clinical prognosis were discovered.

Science, this issue p. eaav1898; see also p. 401

ARTICLE TOOLS

<http://science.sciencemag.org/content/362/6413/eaav1898>

SUPPLEMENTARY MATERIALS

<http://science.sciencemag.org/content/suppl/2018/10/24/362.6413.eaav1898.DC1>

RELATED CONTENT

<http://science.sciencemag.org/content/sci/362/6413/401.full>
<http://stm.sciencemag.org/content/scitransmed/4/156/156ra140.full>
<http://stm.sciencemag.org/content/scitransmed/5/209/209ra153.full>
<http://stm.sciencemag.org/content/scitransmed/7/283/283ra54.full>
<http://stm.sciencemag.org/content/scitransmed/2/38/38ra47.full>

REFERENCES

This article cites 78 articles, 17 of which you can access for free
<http://science.sciencemag.org/content/362/6413/eaav1898#BIBL>

PERMISSIONS

<http://www.sciencemag.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of Service](#)